

Hole doping and weak quenched disorder effects on the 1D Kondo lattice, for ferromagnetic Kondo couplings

Karyn Le Hur

Laboratoire de Physique des Solides, Université Paris-Sud, Bât. 510, 91405 Orsay, France

We investigate the one-dimensional Kondo lattice model (1D KLM) for ferromagnetic Kondo couplings, by using the bosonization method. The ferromagnetic 2-leg spin ladder and the S=1 antiferromagnet occur as new one-dimensional Kondo insulators, for a half-filled band. First, a very small hole-doping makes the charge sector massless and it can produce, either an incommensurate RKKY interaction or a S=1 ferromagnet according to the strength of the ferromagnetic Kondo coupling. Second, we investigate, the effects of a weak and quenched disorder on these two Kondo insulators, by applying renormalization group methods: the Anderson localization is suppressed only in the context of a strong ferromagnetic Kondo coupling.

PACS NUMBERS: 71.27 +a, 75.30 Mb, 75.20 Hr, 75.10 Jm

The Kondo lattice model (KLM) consists of conduction electrons magnetically coupled to a spin array via the Kondo interaction. This model is ruled by two important parameters, namely the hopping strength t of the conduction electrons and the Kondo coupling λ_k , which can be ferromagnetic or antiferromagnetic. Many various phases are expected in the phase diagram, according to the ratio λ_k/t . For instance, the compounds which yield a large and positive ratio λ_k/t are called the “ancestral” Kondo insulators, and they behave, at low-temperatures, like semi-conductors with very small gaps [1]. Conversely, the compounds which yield a small and positive ratio λ_k/t have generally a magnetic ground state. New interesting fixed points of the KLM, have also been investigated, for some years ago, like the heavy-fermion one, which corresponds to an intermediate (positive) ratio λ_k/t [2]: these heavy-fermion compounds are Fermi liquid systems on the brink of magnetism. Exotically, the low-dimensional systems such as the S=1 antiferromagnet [3,4] or the 2-leg spin-ladder [5] have also attracted both theorists and experimentalists: curiously, these low-dimensional spin systems yield a similar ground state and for instance, they are known to develop a spin gap in their singlet-triplet excitations.

Finally, in this Letter, we investigate the case of the 1-dimensional KLM (1D KLM) but originally for a *ferromagnetic* Kondo coupling. In this context, the ferromagnetic 2-leg spin ladder and the S=1 antiferromagnet occur as new interesting Kondo insulators. Explicitly, in this paper, by applying standard bosonization methods [6,7], we begin to yield clearly these original phases of the 1D KLM, for a half-filled band. Then, we view the effects, on these two “similar” Kondo insulators, of a small hole-doping and a weak and dense randomness, by using the well-known replica trick and renormalization-group analysis.

We, first, consider the pure 1d KLM:

$$\mathcal{H} = -t \sum_{i,\sigma} c_{i,\sigma}^\dagger c_{i+1,\sigma} + U \sum_{i,\sigma} n_{i,\sigma} n_{i,-\sigma} + \lambda_k \sum_i \vec{S}_{c,i} \cdot \vec{S}_{f,i} + \lambda_f \sum_i \vec{S}_{f,i} \cdot \vec{S}_{f,i+1} \quad (1)$$

with: ($\lambda_k < 0$, $\lambda_f > 0$, $U > 0$).

Here, $c_{i,\sigma}^\dagger$ ($c_{i,\sigma}$) is the creation (annihilation) operator of conduction electrons, $\vec{S}_c = c_{i,\alpha}^\dagger \frac{\vec{\sigma}_{\alpha\beta}}{2} c_{i,\beta}$ with the Pauli matrix σ and similarly, we introduce $\vec{S}_f = f_{i,\alpha}^\dagger \frac{\vec{\sigma}_{\alpha\beta}}{2} f_{i,\beta}$. The term λ_k describes the ferromagnetic Kondo coupling. An *ad hoc* coupling λ_f between the localized spins is introduced and it will be fixed by physical arguments. A Hubbard interaction between electrons has also been introduced. We approximate that the lattice step $a \rightarrow 0$ and we use the following conventions on the Minkowskian space. The relativistic fermions $c_\sigma(x)$ are separated in left-movers, $c_{L\sigma}(x)$ and right-movers, $c_{R\sigma}(x)$ on the v -cone. By using the bosonization method, the charge $J_{c,L} = c_{L\alpha}^\dagger c_{L\sigma}(x)$ and spin current $\vec{J}_{c,L} = c_{L\alpha}^\dagger \frac{\vec{\sigma}_{\alpha\beta}}{2} c_{L\beta}$ operators can be respectively depicted by a massless scalar field Φ_c^c and a SU(2) matrix g [8]; we just remember the identifications: $J_{c,R} + J_{c,L} = \frac{1}{\sqrt{\pi}} \partial_x \Phi_c^c$ and $J_{c,R} - J_{c,L} = -\frac{1}{\sqrt{\pi}} \Pi_c^c$, where Π_c^c is the moment conjugate to the field Φ_c^c . Then, we decompose the spin operator, into its $k=0$ and $k=\pi$ components as follow [8]:

$$\vec{S}_c \simeq \vec{J}_{c,L}(x) + \vec{J}_{c,R}(x) + cst \exp(i2k_F x) tr(g \cdot \vec{\sigma}) \cos(\sqrt{2\pi} \Phi_c^c) \quad (2)$$

Concerning the localized spins, $\exp(i\sqrt{2\pi} \Phi_f^c)$ is simply replaced by its expectation value; finally, one obtains the general Hamiltonian:

$$\mathcal{H} = H_c + H_s + H_k \quad (3)$$

with:

$$H_c = \int dx \frac{u_\rho}{2K_\rho} : (\partial_x \Phi_c^c)^2 : + \frac{u_\rho K_\rho}{2} : (\Pi_c^c)^2 : \quad (4)$$

$$\begin{aligned}
& + cst.g_3 \cos(\sqrt{8\pi}\Phi_c^c) \\
H_s &= \frac{2\pi v}{3} \int dx : \vec{J}_{c,L}(x) \vec{J}_{c,L}(x) : \\
& + \frac{2\pi v_f}{3} \int dx : \vec{J}_{f,L}(x) \vec{J}_{f,L}(x) : + L \rightarrow R \\
H_k &= \int dx \lambda_1 \vec{J}_{c,L}(x) \vec{J}_{f,L}(x) + \lambda_2 \vec{J}_{c,L}(x) \vec{J}_{f,R}(x) \\
& + L \leftrightarrow R + \lambda_3 tr(g^\dagger \cdot \vec{\sigma}) tr(f \cdot \vec{\sigma}) \cos(\sqrt{2\pi}\Phi_c^c)
\end{aligned}$$

The coupling $g_3 \simeq U/t$ generates the usual $4k_F$ -Umklapp process and the u_ρ , K_ρ parameters of the Tomonaga-Luttinger (TL) liquid [9], are given by:

$$u_\rho K_\rho = v \quad \text{and} \quad \frac{u_\rho}{K_\rho} = v + U/\pi \quad (5)$$

The *dimensionless* parameters $\lambda_{i=1,2,3}$ vary like $(a\lambda_k)$ and the two velocities v_f and v are respectively of order $(a.\lambda_f)$ and $(a.t)$. Due to the $SU(2)$ spin symmetry of electrons, we remember that the spinon velocity is simply equal to v .

The terms λ_1 and λ_2 are *marginal*: the term λ_1 just renormalizes the velocity of spinons; it can be ignored. Conversely, the term λ_2 generates the ferromagnetic Kondo process and the term λ_3 describes the coupling between the staggered magnetizations.

When $a|\lambda_k| \ll v$ (t is large), the term λ_2 is small and it is not really essential at half-filling: it can be forgotten. Conversely, the term λ_3 polarizes the S_f -spins and then, we can take $v_f \simeq v$. The $2k_F$ oscillation becomes commensurate with the alternating localized spin operator. The term λ_3 has the dimension $3/2$ (in the limit $U \ll \lambda_k$) and it opens a mass gap $M \propto \lambda_3^2$, for the charge and all the spin excitations. First, the Umklapp process does not change the physics and it can be dropped out. Second, all the spin excitations are gapped because the spin Hamiltonian is not *conformally invariant*, when we couple 2 $SU(2)_{k=1}$ algebras. To briefly prove that, we note $K = \lambda_3 \langle \cos(\sqrt{2\pi}\Phi_c^c) \rangle$ which is proportional to $-(a\lambda_k)^2$; then, by using the representation of $\vec{J}_{f,(L,R)}$ in terms of the matrix f [8], we obtain the *standard* 2-leg spin-ladder Hamiltonian [10]:

$$H^* = H_s + H_k = W(f) + W(g) + K tr(g^\dagger \cdot \vec{\sigma}) tr(f \cdot \vec{\sigma}) \quad (6)$$

with the well-known $SU(2)_{k=1}$ Wess-Zumino-Witten Hamiltonian [11,12]:

$$\begin{aligned}
W(f) &= \frac{1}{4u^2} \int d^2x tr(\partial_\mu f \partial_\mu f^\dagger) \\
& + \frac{k}{24\pi} \int d^3x \epsilon^{\mu\nu\lambda} tr(f^\dagger \partial_\mu f f^\dagger \partial_\nu f f^\dagger \partial_\lambda f)
\end{aligned} \quad (7)$$

We take $v = 1$ in the definition of $W(f)$ and the central charge $k=1$ is not subject to renormalization. Now, by substituting $g^\dagger = \alpha f^\dagger$, we find that the α -field action

concerns the high energy physics and that the asymptotic behavior of the system is then well-described by the action $2W(f)$ [10,13]: this effective action is not conformally invariant because it is described by a $k=0$ central charge. Consequently, the u -coupling becomes asymptotically free in the Infra-Red limit and it obeys:

$$\beta(u) = \frac{du^2}{d\ln D} = -\frac{1}{4\pi} u^4 \quad (8)$$

By integrating this equation, we find a ground state which looks like a *short-range* Resonating Valence Bond (RVB) system [14] confined inside the characteristic length $\xi \propto \frac{u^2}{D} \exp(\frac{4\pi}{u^2})$ and D is an ultraviolet cut-off. Another consequence is that spin-spin correlation functions decay exponentially or equivalently, that the magnetism is generated by the massive singlet-triplet excitations: the corresponding mass or more commonly, the Haldane gap $\Delta_S \propto |K|/2$ is well-described by the instantonic term $tr g^\dagger \cdot tr f \simeq (tr f)^2$ [13] and we can consider that $D \propto \lambda_3^2$.

When $a|\lambda_k| > v$ (t is small), the electrons are trapped by the lattice spins and the spin system looks like a $S=1$ antiferromagnet. For a strong ferromagnetic Kondo coupling, the $SU(2)$ spin symmetry of electrons is explicitly broken onto $U(1) \times \mathcal{Z}_2$ and the $\mathcal{Z}_2$ symmetry which corresponds to $\sigma_z \rightarrow -\sigma_z$ is spontaneously broken. The term λ_2 is then decomposed into $(\lambda_{2,+}, \lambda_{2,z})$ and the Kosterlitz-Thouless analysis predicts that a mass turns on exponentially with anisotropy:

$$m \propto \lambda_3^2 \exp(-cst/\sqrt{|\Delta J_z|}) \quad (9)$$

with:

$$\lambda_{2,+} - \lambda_{2,z} = |\Delta J_z| \ll 0 \quad (10)$$

Remarkably, in that case, it is well-known [3] that the *same* original RVB ground state occurs and finally, the preceding “weak λ_k -coupling” approach still gives good qualitative results in that case: the $S=1$ antiferromagnet is also ruled by a central charge $k=0$ and also by the term λ_3 . Finally, the mass m of eq. (9) just gives some little details about the singlet-triplet excitations (now the massive triplet excitations are reduced to $S^z = \pm 1$). In conclusion, by decreasing the hopping strength of electrons, we do not hope many drastic changes for a half-filled band, when the Kondo coupling is ferromagnetic.

Even for a *small* hole-doping, the oscillant term λ_3 can be dropped out; then, the system behaves like a metallic state and it is not well-described by the preceding short-range RVB ground state, whatever the ratio $|\lambda_k|/t$. In fact, the Haldane gap is destroyed by the commensurate-incommensurate transition and we can conclude that, conversely to the standard coupled spin-chain problem, the spin sector is not well-described by *any* RVB ground state away from half-filling. Then, according to the ratio

$|\lambda_k|/t$, we obtain the following ground states.

In the regime $a|\lambda_k| \ll v$, the *ad hoc* λ_f -coupling is not well-suited to the situation and we start with $v_f = 0$. The spin system is simply described by a $SU(2)_{k=1}$ algebra but now, we can not omit the small Kondo coupling although it is not subject to renormalization ($\lambda_2^* \simeq \lambda_2 \ll v$). Then, the $SU(2)_{k=1}$ algebra is explicitly described by the spin field: $\vec{J}'_L \simeq \alpha \vec{J}_{c,L} + \beta \vec{J}_{f,R}$ and this formula is correct while $\beta \ll \alpha$. The parameters, α and β are given by the two following constraints:

$$\beta^2 + \alpha^2 = k = 1 \quad (11)$$

and:

$$\beta/\alpha = 3\lambda_2^*/4\pi v \quad (12)$$

Then, by solving these equations we find that only a very small ratio of the localized spins, $\beta^2 = (1 + \frac{16\pi^2 v^2}{9\lambda_2^{*2}})^{-1} \ll 1$ becomes itinerant. Then, the Kondo coupling just renormalizes the velocity of spinon to $v^* = v/\alpha^2 = v \cdot (1 + \frac{9\lambda_2^{*2}}{16\pi^2 v^2})$. Finally, the S_f -spins remain localized and the electrons just polarize them by hopping. By using general results on the TL liquid, we can predict power-law for the spin-spin correlation functions:

$$\langle \vec{J}_f(0,0) \cdot \vec{J}_f(x,0) \rangle \sim \langle \vec{J}_c(0,0) \cdot \vec{J}_c(x,0) \rangle \propto \frac{\cos(2k_F x)}{x^{1+K_\rho}} \quad (13)$$

The system remains antiferromagnetic but we predict the disappearance of the Haldane gap and the enhancement of the RKKY interaction.

When $a|\lambda_k| > v$, the electrons are trapped by spins and the spin system is now composed prevalently of $S=1$ spins and of some rare spinons. For a *very* small hole-doping, a $SU(2)_{k=2}$ algebra is hoped to still describe the ground state and we can not use a “weak λ_k -coupling” approach in that case. Anyway, the ground state is quite easy to intute. Indeed, since $\lambda_3 = 0$, we do not expect any $k = \pi$ fermionic excitations and finally, this implies that the $S=1$ spins form a ferromagnetic ground state: this is also in agreement with a metallic ground state.

Finally, we expect a cross-over from an antiferromagnetic and incommensurate RKKY situation towards a ferromagnetic ground state (formed by the $S=1$ spins) by decreasing the hopping strength t of electrons.

Now, for a half-filled band, we investigate the effects of a *small and dense* quenched disorder on the two original Kondo insulators, namely the ferromagnetic 2-leg spin ladder and the $S=1$ antiferromagnet. We use an Abelian representation of the $SU(2)$ matrices f and g and by introducing the usual boson fields $\Phi_\pm = (\Phi_c \pm \Phi_f)/\sqrt{2}$ for the spin sector and their canonical conjugate momenta $\Pi_\pm = \partial_x \tilde{\Phi}_\pm$, we obtain:

$$H^* = \sum_{\nu=+,-} \int dx \frac{u_\nu}{2K_\nu} : (\partial_x \Phi_\nu)^2 : + \frac{u_\nu K_\nu}{2} : (\Pi_\nu)^2 : \quad (14)$$

$$+ \int dx \frac{cst}{\alpha^2} [\lambda_4 \cos(\sqrt{4\pi}\Phi_-) + 2\lambda_5 \cos(\sqrt{4\pi}\tilde{\Phi}_-) - \lambda_6 \cos(\sqrt{4\pi}\Phi_+)] \cos(\sqrt{2\pi}\Phi_c^c)$$

As seen before, the Kondo term does not change the physics and it has not been introduced; then, the parameter K_- takes into account the fact that the $SU(2)$ spin symmetry is explicitly broken for a strong ferromagnetic Kondo coupling: in that case K_- is large and the massive excitations are reduced to the doublet $S^z = \pm 1$ well-described by the term $\cos(\sqrt{4\pi}\phi_+)$ [10]. Finally, we want to remember that a “weak λ_k -coupling” approach still gives good results concerning the $S=1$ antiferromagnet ground state.

Here, we apply renormalization group methods, first used by Giamarchi and Schulz [15] (in the context of randomness in a TL liquid). We introduce the complex random impurity potential,

$$H_{imp} = \sum_\sigma \int dx \xi(x) c_{\sigma L}^\dagger c_{\sigma R} + hc \quad (15)$$

with the Gaussian distribution:

$$P_\xi = \exp(-D_\xi^{-1} \int dx \xi^\dagger(x) \xi(x)) \quad (16)$$

We can omit forward scatterings, because the $k=0$ random potential just renormalizes the chemical potential and it does not affect the fixed point properties. In order to deal with the quenched disorder, we use the well-known *replica* trick [16]. Due to the definition of $\xi(x)$, we are limited to a weak randomness treatment and we incorporate only the first contribution (order) in D_ξ . Then, there is no coupling between different replica indices, which will be omitted below. By applying a standard renormalization group analysis up to the lowest order in λ_k and D_ξ , we obtain the complete flow:

$$\frac{dD_\xi}{dl} = (3 - \frac{K_+}{2} - \frac{K_-}{2} - K_\rho) D_\xi \quad (17)$$

$$\frac{d\lambda_4}{dl} = (2 - K_- - \frac{K_\rho}{2}) \lambda_4 \quad (18)$$

$$\frac{d\lambda_5}{dl} = (2 - \frac{1}{K_-} - \frac{K_\rho}{2}) \lambda_5 \quad (19)$$

$$\frac{d\lambda_6}{dl} = (2 - K_+ - \frac{K_\rho}{2}) \lambda_6 \quad (20)$$

$$\frac{dK_+}{dl} = -\frac{1}{2}(D_+ + \lambda_6^2) K_+^2 \quad (21)$$

$$\frac{dK_-}{dl} = -\frac{1}{2}(D_- + \lambda_4^2) K_-^2 + 2\lambda_5^2 K_-^2 \quad (22)$$

$$\frac{dK_\rho}{dl} = -\frac{1}{2}u_\rho \left[\left(\frac{D_-}{u_-} + \frac{D_+}{u_+} \right) + \left(\frac{\lambda_4^2 + 4\lambda_5^2}{u_-} + \frac{\lambda_6^2}{u_+} \right) \right] K_\rho^2 \quad (23)$$

with the notations: $D_\nu = 2 \frac{D_\xi \alpha}{\pi u_\nu^2} [\frac{u_\nu}{u_\rho}] K_\rho$ and $\tilde{\lambda}_\nu = \frac{\lambda_\nu}{2\pi u_\nu}$ and where we did not display the irrelevant equations

to the following discussions. We can particularly notice that in this problem, there is no term like $D_\xi \lambda_{(4,6)}$ generated by perturbation. Finally, the fixed point properties clearly depend on the initial conditions, on the pure system.

For instance, if we consider the 2-leg spin ladder situation, we initially have $2 - K_- - \frac{K_\rho}{2} > 0$, $2 - \frac{1}{K_-} - \frac{K_\rho}{2} > 0$ and $2 - K_+ - \frac{K_\rho}{2} > 0$. By using the above equations, we can easily conclude that, in this case, D_ξ scales to the strong coupling regime before the couplings λ_ν : the Haldane gap vanishes and K_ρ tends to zero. Although in the strong coupling regime, our treatment for randomness breaks down, we can guess the fixed point properties. Indeed, the charge gap $\Delta_c \propto \lambda_k^2$ is small and it seems plausible to expect that the TL liquid will be (still) quenched by a strong disorder, to give a transition into the famous Anderson localization state [17].

Now, if we branch a strong ferromagnetic Kondo coupling by decreasing the hopping strength t , we initially have $2 - K_- - \frac{K_\rho}{2} < 0$, $2 - \frac{1}{K_-} - \frac{K_\rho}{2} > 0$ and $2 - K_+ - \frac{K_\rho}{2} > 0$ and $K_- \gg 1$ since the SU(2) spin symmetry is explicitly broken onto $U(1) \times \mathcal{Z}_2$. By using the eqs. (21), (23), we can clearly see that K_+ and K_ρ are also reduced by the randomness and gaps open in Φ_+ and Φ_-^c : finally, $K_- \rightarrow +\infty$ at the fixed point and the disorder is reduced to zero. The coupling λ_6 scales to the strong coupling limit and we can check that the Haldane gap is then given by the single instantonic term $\cos(\sqrt{4\pi}\phi_+)$. Then, the Haldane gap is maintained and it is not surprising if we remember that the charge gap $\Delta_c \propto \lambda_k^2$ is large in the pure system and finally it is even larger than the Haldane gap ($\Delta_S \propto 1/2 \cdot \lambda_k^2$): the electrons are quite quenched by magnetic impurities and they are not sensitive to a small quenched disorder.

Summarizing, for a half-filled band, the 2-leg spin ladder and the S=1 antiferromagnet occur as new original Kondo insulators; both, they yield a short-range RVB ground state and massive singlet-triplet excitations. But, these two similar Kondo insulators do not equivalently react by hardly hole-doping or by applying a quenched and dense disorder. First, a strong ferromagnetic Kondo coupling is essential to maintain the Haldane gap and to prevent a transition into an Anderson localization state. However, we expect the step of a charge localization in the transport properties, whatever the ratio λ_k/t : localization by the “randomness” in the spin-ladder context and localization by the spins in the case of a strong ferromagnetic Kondo coupling. Second, by hardly hole-doping, the ground state becomes very dependent on the parameter $|\lambda_k|/t$. Indeed, for a ratio $|\lambda_k|/t \ll 1$, the antiferromagnetic correlations are enhanced due to the emergence of an incommensurate RKKY interaction. Conversely, when the ferromagnetic Kondo coupling is large or $|\lambda_k|/t > 1$, we expect another curious change by hardly hole-doping: the residual spin system, preva-

lently formed by S=1 spins yields a ferromagnetic ground state: it minimizes the kinetic energy. Finally, if we remove a S_f -spin of the lattice instead of an electron, we predict the same conclusions concerning the spin sector, since $\lambda_3 = 0$ in that case too, but the system remains insulator because the spins can not move.

-
- [1] G.Aeppli and Z. Fisk, Comments Cond. Matt. Phys. **16** (1992) 155.
 - [2] B. Coqblin, J. Arispe, J. R. Iglesias, C. Lacroix and Karyn Le Hur, J. Phys. Soc. Jpn **65** (1996) Suppl. B 64.
 - [3] Theoretically: F.D.M. Haldane, Phys. Lett **93A** (1983) 464.
 - [4] Experimentally: J.P. Renard, M. Verdguer, L.P. Regnault, W.A.C. Erkelens, J. Rossat-Mignot and W.G. Stirling, Europhys. Lett **3** (1987) 945.
 - [5] For a theoretical introduction to spin-ladders: E. Dagotto, J. Riera and D.J. Scalapino, Phys. Rev. B **45** (1992) 5744; M. Sigrist, T.M. Rice and F.C. Zhang, Phys. Rev. B **49** (1994) 12058; S.R. White, R.M. Noack and D.J. Scalapino, Phys. Rev. Lett. **73** (1994) 886; E. Dagotto and T.M. Rice, Science **271** (1996) 618.
 - [6] See, e.g., V.J. Emery, in *Highly Conducting One-Dimensional Solids*, edited by J.T. Devreese, R.P. Evrard and V.E. Van.Doren, (Plenum, New York, 1979), p.327; J. Solyom, Adv.Phys. **28** (1979) 201; S. Sachdev and R. Shankar, Phys. Phys. Rev. B **38** (1988) 826.
 - [7] Concerning the KLM, we suggest: S.P. Strong and A.J. Millis, Phys. Rev. Lett. **69** (1992) 2419; S. Fujimoto and N. Kawakami, J. Phys. Soc. Jpn **63** (1994) 4322; O. Zachar, S.A. Kivelson and V.J. Emery, Phys. Rev. Lett. **77** (1996) 1342; S.R. White and I. Affleck, Phys. Phys. Rev. B **54** (1996) 9862.
 - [8] I. Affleck in *Fields, Strings and Critical phenomena* edited by E. Brezin and J. Zinn-Justin (North-Holland, Amsterdam, 1990).
 - [9] J. M Luttinger, J. Math. Phys. **4** (1963) 1154, H.J. Schulz in *Mesoscopic Quantum Physics, Les Houches, Session LXI, 1994*, edited by E. Akkermans, G. Montambaux, J.L. Pichard and J. Zinn-Justin (Elsevier, Amsterdam, 1995), p. 533.
 - [10] D.G. Shelton, A.A. Nersesyan and A.M. Tsvelik, Phys. Rev. B **53** (1996) 8521.
 - [11] V.G. Knizhnik and A.B. Zamolodchikov, Nucl. Phys. B **247** (1984) 83.
 - [12] E. Witten, Commun. Math. Phys. **92** (1984) 455-47.
 - [13] Karyn le Hur, (unpublished).
 - [14] P.W. Anderson, Science **235** (1987) 1196.
 - [15] T. Giamarchi and H.J. Schulz, Phys. Rev. B **37** (1988) 325.
 - [16] S.F. Edwards and P.W. Anderson, J. Phys. F **5** (1975) 965.
 - [17] P.W. Anderson, Phys. Rev. **109** (1958) 1492.